



Cadilac

800 WP



Fungicide for control of grey mould and downy mildew on roses and early and late blights on potatoes and tomatoes, leaf spots and anthracnose on onions and french beans (Dawa ya kuzuia grey mold na downy mildew kwenye waridi, early blight na late blight kwenye viazi na nyanya na leaf spots na anthracnose kwenye vitunguu na michiri)

GUARANTEE (DHAMANA):
Mancozeb 800g/kg.

COMMERCIAL & AGRICULTURAL CLASS
(KUNDI LA BIAISHARA NA KILIMO)
REGISTRATION NO. (NAMBARI YA USAJILI): PCPB (R) 1105

READ THE LABEL BEFORE USE
(SOMA KIBANDIKO KABLA YA KUTUMIA)
KEEP LOCKED OUT OF REACH OF CHILDREN
(FUNGIA MBALI NA WATOTO)

Batch No. (Nambari ya kifurushi) **SEE PACKING**
Date of Manufacture
(Tarehe ya kutengenezwa): **SEE PACKING**

In case of emergency, call (24hr) toll free number, (Wakati wa dharura, piga simu (masaa 24) bila malipo kwa): 0800720021 / 0800730030

Cadilac® is a Trademark of (alama ya biashara ya):
Greenlife Crop Protection Africa Ltd

Distributed by (Wasambazaji):



P.O. Box 24842 - 00100, Nairobi, Kenya.
Tel: +254 20 244120/12.
Mob: +254 735 545644 +254 722 563698.
+254 738 980267
info@greenlife.co.ke | www.greenlife.co.ke



Net Contents (Kiasi)
10 x 1kg



Manufacturer (Watengenezaji):
AGRIA SA, BULGARIA
Shelf Life (Maisha Rafuini): 2 years from the date of manufacture when stored in unopened container in a cool dry place. (Mikaa miwili baada ya tarehe ya kutengenezwa kama imehifadhiwa kwenye chombo chake asili)

DESCRIPTION: CADILAC 800 WP: is a contact and protective fungicide for control of grey mould and downy mildew on roses and early and late blights on potatoes and tomatoes, leaf spots and anthracnose on onions and french beans. It has protective action with a good long-lasting effect of about 15 days.

MODE OF ACTION: Mancozeb is a non-systemic fungicide with protective and contact action. It acts on multiple sites in fungal cells by disrupting lipid metabolism, respiration and production.

APPLICATION RATES

| Crop | Disease | Rate |
|----------------------|-----------------------------|-------------------------------------------------|
| Roses | Grey mould and downy mildew | 2 kg per hectare in 1000 litres of spray volume |
| Potatoes, Tomatoes | Early and late blights | (40g/20L water) |
| Onions, French beans | Anthracnose and leaf spots | |

PRE HARVEST INTERVAL: 7 days in potato and tomato, 14 days in onions and 21 days in french beans.

APPLICATION TIMING: CADILAC 800 WP can be used during all the vegetative periods of the crops 2-4 times. It is applied before or after infection, at 10-14 day intervals

PREPARATION OF SPRAY MIXTURE: Fill spray tank with half the amount of required water. Add required amount of CADILAC 800 WP and fill the water to the required level. Agitate to ensure thorough mixing.

RE-ENTRY PERIOD: Do not enter treated areas before 24 hours after application unless wearing appropriate protective clothing when necessary.

PHYTOTOXICITY: CADILAC 800 WP is not Phytotoxic when used as recommended.

SAFETY PRECAUTIONS: Avoid contact with mouth, skin and eyes. When using, do not eat, drink or smoke. Wear overalls, safety glasses and impervious gloves. Wash contaminated clothing and other protective equipment before storage or re-use.

STORAGE: Store in the closed, original container in a cool, well-ventilated area. Do NOT store for prolonged periods in direct sunlight. Keep out of the reach of children.

DISPOSAL: Rinse containers before disposal. Puncture or crush empty containers and dispose in a safe designated place away from water sources. Do not reuse empty containers.

ENVIRONMENTAL HAZARDS: CADILAC 800 WP is slightly toxic to fish and birds. Do not contaminate streams, rivers or waterways with the chemical or used containers.

FIRST AID TREATMENT:
Inhalation: If inhaled, move to fresh air and keep at rest. Obtain medical advice if symptoms persist.
Skin contact: Wash off immediately with soap and plenty of water. Seek medical aid if symptoms persist.
Eye contact: Immediately rinse eyes with plenty of water and seek medical advice.
Ingestion: Move victim to fresh air. Keep patient at rest and seek medical advice.

TOXICOLOGICAL INFORMATION:
Product is poisonous if swallowed or absorbed by skin contact.
Symptoms of poisoning: Nausea, vomiting, diarrhoea, headache, ataxia, confusion and fatigue.
Antidote: No known antidote.
Note to the physician: Treatment should be symptomatic.

In case of emergency, call (24hr) toll free number 0800720021 or 0800730030

NOTICE TO USERS: This pest control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions.

WARRANTY: Seller's guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition. The user bears the risk for damage resulting from factors beyond the manufacturers control

MAELEZO: CADILAC 800 WP ni dawa ya mguso ya kuzuia grey mould na downy mildew kwenye waridi, early blight na late blight kwenye viazi na nyanya na leaf spots na anthracnose kwenye vitunguu na michiri. Inaanza magojwa na husalia kwenye mimea kwa muda wa hadi siku 15.

INAVUFANYA KAZI: Mancozeb hufanya kazi kwa kuzuia ukauji wa vitini vya vyobabishi magojwa na kwa mguso. Huatafika mawasiliano muhimu kwenye sehemu mbalimbali kwa chembechembe za fungi ambazo huhitafika ueneaji na ukauji.

KIPIMO CHA KUTUMIA

| Mimea | Magonjwa | Kipimo |
|-------------------|-----------------------------|------------------------------------------|
| Waridi | Grey mould na downy mildew | Kilo 2 kwa hektari kwa lita 1000 ya maji |
| Viazi, nyanya | Early blight na late blight | (Gramu 40 kwa lita 20 ya maji) |
| Vitunguu, michiri | Leaf spots na anthracnose | |

MUDA SALAMA WA KUSUBIRI KABLA YA KUVUNA: Siku 7 kwa nyanya na viazi, siku 14 kwa vitunguu na siku 21 kwa michiri.

WAKATI WA KUTUMIA: CADILAC 800 WP yaweza kutumia wakati wovote katika maisha ya mmea. Tumia mara 2 hadi 4 katika mmea mmoja. Tumia kabla au baada ya kuonekana kwa ugonjwa na urudie kila baada ya siku 10 hadi 14.

JINSI YA KUCHANGANWA: Tia maji kwenye bomba hadi nusu halafu onyesha kiwango cha CADILAC 800 WP kinachohitajiwa. Kisha ongeza maji yaliyobakia huku ukikogorora vizuri.

MUDA SALAMA WA KUINGIA PALIPONYUNZIWA: Uingie paliponyunziwa dawa hihi kabla ya muda wa masaa 24 kukamilika. Kiwa utazamikia kuingia, hakikisha uwevava mavazi rasmi ya kujikinga.

TAHADHARI: Epuka dawa kuingia kwenye mdomo, macho, au kugusana na ngozi. Usile, kunywa wala kuvuta sigara unapoitumia dawa hihi. Vaa mavazi ya kinga k.v. kinga ya uso, mifuko ya raba ya mikono, na overall unapochanganyia na kunyunyiza dawa hihi. Baada ya kutumia dawa, osha vizuri mavazi ya kujikinga.

KUHIFADHI: Dawa hihi yafaa kuhifadhiwa kwenye chombo chake asili, pahali pakavu, pasipo na joto jingi huku ukiepusha kugusana na vyaku. Weka mbali na watoto.

UTUPAJI: Chombo kitupu cha dawa hihi chafaa kusafisha mara tatu, kutobolewa na kutupwa au kuzikwa pahali palipoitungwa, mbali na chemchemi au hifadhi ya maji. Ustumie chombo kitupu kwa kazi zingine zozote.

ADHARI KWA MZINGIRA: CADILAC 800 WP yaweza kudharu samaki na ndege. Kwa hivyo dawa yenyeve, mabaki yake au chombo kitupu hazifai kutumia au kutupwa kwa njia itakayochafua visima, mito, vidimbi, au chemchemi ya maji.

HUDUMA YA KWANZA:
Ikrutiva kwa mapua: Eleka mhadhiriwa palipo na hewa safi na unutilize. Tafuta ushauri wa daktari.
Ikgusana na ngozifisha: ngozizi vizuri kwa maji mengi na sabuni. Tafuta ushauri kwa daktari.

Ikingia kwa macho: Fungua na kuyaosha macho kwa maji mengi na utafute ushauri wa daktari.
Ikimwaa: Peleka mhadhiriwa palipo na hewa safi. Mutlize na utafute ushauri wa daktari.

MAELEZO KUHUSU SUMU: Inadhuru ikimezwa au ikigusana na ngozi. Dalili za kusumika: Kuhisi kutapika au kutapika, kuhara, kuuma na kitwaa, kukosa mwelekeo, kuchanganikiwa, kuhisi uchowa mwilini.

Makata: Hakuna tiba maalum.
Arifa kwa daktari: Tibu kulingana na dalili za kusumika.

MALEZELO KUHUSU SUMU: Inadhuru ikimezwa au ikigusana na ngozi. Dalili za kusumika: Kuhisi kutapika au kutapika, kuhara, kuuma na kitwaa, kukosa mwelekeo, kuchanganikiwa, kuhisi uchowa mwilini.

Arifa kwa daktari: Tibu kulingana na dalili za kusumika.

Wakati wa dharura, piga simu (masaa 24) bila malipo kwa 0800720021 / 0800730030

ILANI KWA MTUMIAJI: Dawa hihi sharti itumie kulingana na maagizo yaliyo kwenye kibandiko. Ni hatia chini ya sherha za madawa ya kungumiza waduu waharibiti kutumia au kuhifadhi dawa hihi katika hali itoye salama.

THIBITISHO: Dhama ya muuzaji ni maelezo yaliyo kwenye kibandiko ambayo lazima yafuatue ikamilifu. Mnuuzi lazima akubali hatari zote zinazoweza kutokana na matumizi ya dawa hihi kwa binadamu au maji na lazima akubali dawa katika hali hiyo.

